

# The Relationship Between Depression and Internet Addiction

by Kimberly S. Young and Robert C. Rodgers

Paper published in *CyberPsychology & Behavior*, 1(1), 25-28, 1998

## ABSTRACT

Prior research has utilized the Zung Depression Inventory (ZDI) and found that moderate to severe rates of depression coexist with pathological Internet use.<sup>1</sup> Although the ZDI was utilized for its expediency with on-line administration, its limitations include poor normative data and less frequent clinical use. Therefore, this study utilized the Beck Depression Inventory (BDI), which has more accurate norms and frequent usage among dual diagnostic patient populations. An on-line survey administered on a World Wide Web site utilized the BDI as part of a larger study. A total of 312 surveys was collected with 259 valid profiles from addicted users, which again supported significant levels of depression to be associated with pathological Internet use. This article discusses how a treatment protocol should emphasize the primary psychiatric condition if related to a subsequent impulse control problem such as pathological Internet use. Effective management of psychiatric symptoms may indirectly correct pathological Internet use.

Prior research has identified the existence of addictive Internet use, which has been associated with significant social, psychological, and occupational impairment.<sup>2</sup> Addicts in this study used the Internet an average of 38 hr per week for nonacademic or non-employment purposes, which caused detrimental effects such as poor grade performance among students, discord among couples, and reduced work performance among employees. This is compared to non-addicts who used the Internet an average of 8 hr per week with no significant consequences reported. Predominantly, the interactive capabilities of the Internet such as chat rooms or on-line games were seen to be the most addictive. This type of behavioral impulse control failure, which does not involve an intoxicant, was seen as most akin to pathological gambling. Therefore, a formal term utilized in this article is *pathological Internet use* (PIU) to refer to cases of addictive Internet use.

Research in the addictions field has shown that psychiatric illnesses such as depression are often associated with alcoholism<sup>3</sup> and drug addiction.<sup>4</sup> Further, research has shown that other addictive behaviors overlap with depression—for example, eating disorders<sup>5,6</sup> and pathological gambling.<sup>7-9</sup> Although the concept of Internet addiction has gained credibility among mental health professionals both in academic and clinical realms, little research has been conducted to examine if similar underlying psychiatric illnesses may contribute to such Internet abuse.<sup>1</sup>

Therefore, the objective of this study was to assess depression and compare such results to other established dual diagnostic populations. Young<sup>1</sup> utilized the Zung Depression Inventory<sup>10</sup> (ZDI), which suggested that increased levels of depression are associated with moderate to severe levels of PIU. However, the ZDI yields limited clinical utility; therefore, this study used the Beck Depression Invento#1 (BDI) because it is a more psychometrically and clinically valid

instrument to further investigate the effects of depression on PIU. Finally, this study also attempted to increase its sample size from the previous examination ( $N = 99$ ) to improve generalizability of results.

## Method

**Subjects**  
**Materials**  
**Procedures**

## Results

## Depression and Addiction Discussion

## References

---

## METHOD

### *Subjects*

Subjects were self-selected active Internet users who responded to postings on electronic support groups and those who searched for the keywords *Internet* or *addiction* on popular Web search engines (e.g., Yahoo).

### *Materials*

An on-line survey was constructed for this study. The survey exists as a World Wide Web (WWW) page (located at <http://www.pitt.edu/ksy/survey.html>) implemented on a UNIX-based server that captures the answers into a text file. The on-line survey administered a structured diagnostic questionnaire that modified the *DSM-IV* criteria for pathological gambling<sup>2</sup> to classify subjects as addicted or non-addicted, followed by administration of the BDI, the Sixteen Personality Factor Inventory,<sup>15</sup> and Zuckerman's Sensation Seeking Scale,<sup>13</sup> as part of a larger study. Finally, demographic information was also gathered.

### *Procedures*

The WWW location of the survey was submitted to several popular search engines available to assist on-line users in finding Web pages of interest. On-line users entering keyword searches for *Internet* or *addiction* would find the survey and have the option to follow the link to the survey in order to fill it out. Additionally, a brief description of the study along with the WWW address of the survey was advertised on prominent electronic support groups geared toward Internet addiction (e.g., the Internet Addiction Support Group and the Web-aholics Support Group). Answers to the survey were sent in a text file directly to the principal investigator's electronic

mailbox for analysis. Respondents who answered "yes" to five or more of the criteria were classified as addicted Internet users for inclusion in this study.

## **RESULTS**

A total of 312 surveys were collected, resulting in 259 valid geographically dispersed profiles from addicted users. The sample included 130 males with a mean age of 31 and 129 females with a mean age of 33. Educational background was as follows: 30% had a high school degree or less, 38% had an associate's or bachelor's degree, 10% had a master's degree or doctorate, and 22% were still in school. Of the subjects, 15% had no vocational background (e.g., homemaker or retired), 31% were students, 6% were blue-collar workers (e.g., factor worker or auto mechanic), 22% were non-tech white-collar workers (e.g., school teacher or bank teller), and 26% were high-tech white-collar workers (e.g., computer scientist or systems analyst).

Occupational type appears to be a determinant in the level of Internet usage in this study. These results suggest that non-tech or high-tech white-collar workers are more likely to become addicted to the Internet than are blue-collar workers. White-collar employment may offer wider access to the Internet and greater salary potential, making the purchase of a home computer more affordable compared to those in blue-collar types of employment, which may explain these results.

Results from the BDI were a mean of 11.2 (*SD* 13.9), indicating mild to moderate levels of depression compared to normative data. Prior research showed that analysis of the ZDI provided a mean of 38.56 (*SD* = 10.24), also indicating mild to moderate levels of depression when compared to normal populations.~ Therefore, the BDI yielded similar results as the prior work suggesting that depression is a significant factor in the development of PIU.

## **DEPRESSION AND ADDICTION DISCUSSION**

As noted with other addictive disorders, our findings suggest that increased levels of depression are associated with those who become addicted to the Internet. This suggests that clinical depression is significantly associated with increased levels of personal Internet use. These results should be interpreted with caution, however, as self-selected sample biases exist in this study coupled with the questionable accuracy of on-line responses.

This study suggests that accurate assessment of depression and PIU can improve early detection, especially when one is masked by primary symptoms of the other diagnosis. It is likely that low self-esteem, poor motivation, fear of rejection, and the need for approval associated with depressives contribute to increased Internet use, as prior research indicated that the interactive capabilities available on the Internet were found to be most addictive.<sup>2</sup> It is plausible that depressives are drawn to electronic communication because of the anonymous cover granted to them by talking with others through fictitious handles, which helps them overcome real-life interpersonal difficulties. Kiesler et al.<sup>14</sup> found that computer-mediated communication weakens social influence by the absence of such nonverbal behavior as talking in the head set, speaking loudly, staring, touching, and gesturing. Therefore, the disappearance of facial expression, voice

inflection, and eye contact makes electronic communication less threatening, thereby helping the depressive to overcome the initial awkwardness and intimidation in meeting and speaking with others. This anonymous two-way talk also helps depressives feel comfortable sharing ideas with others thanks to the personal control over the level of their communication, as they have time to plan, contemplate, and edit comments before sending an electronic message. Therefore, the treatment protocol should emphasize the primary psychiatric condition, if related to a subsequent impulse control problem, as addictive Internet use. Effective management of such psychiatric symptoms may indirectly correct PIU.

Based on the findings, it is concluded that evaluation of suspected cases of PIU should include assessment for depression. These results, however, do not clearly indicate whether depression preceded the development of such Internet abuse or if it was a consequence. Young<sup>2</sup> showed that withdrawal from significant real-life relationships is a consequence of PIU. Therefore, the possibility exists that increased levels of social isolation subsequent to excessive time spent in front of a computer may result in increased depression rather than be a cause of such Internet overuse. Therefore, further experimentation with a more comprehensive level of analysis is necessary to examine cause and effect. Data collection should also include patients in treatment to eliminate the methodological limitations of an on-line survey and to improve the clinical utility of the information gathered. Finally, although it is unclear how PIU compares to other established addictions, future research should investigate if clinical depression is an etiologic factor in the development of any addictive syndrome, be it alcohol, gambling, or the Internet.

## REFERENCES

1. Young, K.S. (1997, April 11). *Levels of depression and addiction underlying pathological Internet use*. Poster presented at the annual meeting of the Eastern Psychological Association, Washington, DC.
2. Young, K.S. (1996, August 10). *Internet addiction: The emergence of a new clinical disorder*. Paper presented at the 104th annual meeting of the American Psychological Association, Toronto.
3. Capuzzi, D., & Lecoq, L.L. (1983). Social and personal determination of adolescent use and abuse of alcohol and marijuana. *Personnel and Guidance Journal*, 62, 199-205.
4. Cox, W.M. (1985). Personality correlates of substance abuse. In M. Galizio & S.A. Maisto (Eds.), *Determinants of substance abuse: Biological, psychological, and environmental factors* (pp.209-246). New York: Plenum.
5. Lacey, H.J. (1993). Self-damaging and addictive behavior in bulimia nervosa: A catchment area study. *British Journal of Psychiatry*, 163, 190-194. Lacey, H.J. (1993). Self-damaging and addictive behavior in bulimia nervosa: A catchment area study. *British Journal of Psychiatry*, 163, 190-194.
6. Lesieur, H.R., & Blume, S.B. (1993). Pathological gambling, eating disorders, and the

- psychoactive substance use disorders. *Journal of Addictive Diseases*, 12(3), 89-102.
7. Blaszczynski, A., McConaghy, N., & Frankova, A. (1991). Sensation seeking and pathological gambling. *British Journal of Addiction*, 81, 113-117.
  8. Criffiths, M. (1990). The cognitive psychology of gambling. *Journal of Gambling Studies*, 6, 31~2.
  9. Mabilia, P. (1993). Gambling as a rational addiction. *Journal of Gambling Studies*, 9(2), 121-151.
  10. Zung, W.K. (1965). *Self-rating depression scale*. New York; Springer-Verlag.
  11. Beck, A.T., Ward, C.M., Mendeleon, M., Mock, J.F., & Erbaugh, J.K. (1961). An inventory for measuring depression. *Archives of General Psychiatry*, 4, 5~-571.
  12. American Psychiatric Association. (1994). *Diagnostic and statistical manual of mental disorders* (4th ed.). Washington, DC: Author.
  13. Zuckerman, M. (1979). *Sensation seeking behavior: Beyond the optimal level of arousal*. Hillsdale, NJ: Erlbaum.
  14. Kiesler, S., Siegal, I., & McGuire, T.W. (1984). Social psychological aspects of computer-mediated communication. *American Psychologist*, 39(10), 1123~134.
  14. Kiesler, S., Siegal, I., & McGuire, T.W. (1984). Social psychological aspects of computer-mediated communication. *American Psychologist*, 39(10), 1123~134.
  15. Cattell, R. (1975). *Sixteen Personality Factor Inventory*. The Institute of Personality and Ability, Inc., Champaign, IL.